arXiv:cond-mat/0303601v1 [cond-mat.stat-mech] 28 Mar 2003

An appropriate candidate for exact distribution of closed random walks using quantum groups

S. A. Alavi †, **M. Sarbishaei**

Department of Physics, Ferdowsi University of Mashhad, Mashhad, P. O. Box 1436, Iran

Abstract. *We show that the structure of the quantum group $su_q(2)$ is intimately related to the random walks on a two dimentional lattice. Using this connection we obtain an appropriate candidate for the exact area distribution of closed random walks of length N on a two dimensional square lattice. We compare our results with exact enumeration.*

Key Words. Quantum groups, Random walks.

PACS Numbers: 05.45.+b, 72.15.Qm, 72.10.Bg

1.Introduction. Let us consider a spinless electron on a two dimensional lattice and submitted to a uniform magnetic field along the z-direction and perpendicular to the plane of motion. The Hamiltonian of the system is:

$$h = \frac{1}{2m}(\vec{p} + e\vec{A})^2. \tag{1}$$

The system is not invariant under translations but there is an invariance under the so-called magnetic translation operators [1]:

$$w(\vec{a}) = \exp(\frac{i}{\hbar}\vec{a}.\vec{k}), \tag{2}$$

where $\vec{a}$ is an arbitrary two dimensional lattice vector and $\vec{k}$ is:

$$\vec{k} = k_x\hat{x} + k_y\hat{y} = \vec{p} + e\vec{A} + e\vec{r}\times\vec{B}. \tag{3}$$

There is a relationship between magnetic translation operators, quantum groups, Landau levels and the quantum Hall effect which has been the subject of study in several references [e.g.,2-4]. It is shown that $w(a)$ satisfy the following relation:

$$w(\vec{a})w(\vec{b}) = \exp(i\frac{e}{2\hbar}\vec{B}.(\vec{a}\times\vec{b}))w(\vec{a}+\vec{b}), \tag{4}$$

where $\vec{a}\times\vec{b} = a_1b_2 - a_2b_1$,which is exactly the q-commutation relation:

$$w(\vec{a})w(\vec{b}) = qw(\vec{b})w(\vec{a}), \tag{5}$$

---

†.*Corresponding author.*
E-mail:*Alavi@sttu.ac.ir* and *s_alialavi@hotmail.com*

where $q = \exp(i\frac{e}{\hbar}\vec{B}.(\vec{a}\times\vec{b}))$. We can write q in the form $q = e^{i\gamma}$ where $\gamma = 2\pi\frac{\Phi}{\Phi_0}$. Φ and Φ_0 are the magnetic flux through the unit cell and the quantum of the flux respectively :

$$\Phi = \vec{B}.(\vec{a}\times\vec{b}), \Phi_0 = \frac{h}{e}. \tag{6}$$

Here $\vec{a}$ and $\vec{b}$ are two perpendicular unit vectors which build the lattice unit cell, $\mid a \mid=\mid b \mid= 1$. Using (4) and (5) it is easy to show that the following combinations of the magnetic translation operators :

$$j_+ = \frac{w(\vec{a}) + w(\vec{b})}{q - q^{-1}}, \tag{7}$$

$$j_- = -\frac{w(\vec{-a}) + w(\vec{-b})}{q - q^{-1}}, \tag{8}$$

and $q^{j_3} = w(\vec{b} - \vec{a}) = j_z$ form the $su_q(2)$ algebra. The $su_q(2)$ algebra is the q-deformation of the lie algebra $su(2)$ [5-8], with generators satisfying the commutation relations:

$$[j_3, j_\pm] = \pm j_\pm. \tag{9}$$

$$[j_+, j_-] = \frac{q^{2j_3} - q^{-2j_3}}{q - q^{-1}}. \tag{10}$$

We observe that (4), (5) and the $su_q(2)$ algebra are invariant under following transformations:

$$w(\vec{a}) \to g(q)w(\vec{a}), \tag{11}$$

$$w(-\vec{a}) \to g^{-1}(q)w(-\vec{a}), \tag{12}$$

or equivalently:

$$j_+ \to g(q)j_+, \tag{13}$$

$$j_- \to g^{-1}(q)j_-, \tag{14}$$

$$j_z \to j_z, \tag{15}$$

where $g(q)$ is an arbitrary function of q.

2. Areas distribution of closed random walks.

Bellissard et al [9] derived the distribution of areas of closed random walks using noncommutative geometry. Their method was based on the Harper model. The Harper model Hamiltonian [10] is given by:

$$H = \sum_{|a|=1} w(\vec{a}) = w(\vec{a}) + w(\vec{b}) + w(-\vec{a}) + w(-\vec{b}). \tag{16}$$

We define the trace per unit area such that

$$T(w(\vec{m}_1)w(\vec{m}_2)) = \delta_{\vec{m}_1+\vec{m}_2,0}. \tag{17}$$

Using (16) and (17) it is easy to see that :

$$T(H^N) = \sum_{\Gamma} e^{\frac{i\gamma}{2}A(\Gamma)}, \tag{18}$$

where the sum is over the set of closed paths starting at the origin of length N, and $A(\Gamma)$ is the algebraic area closed by Γ. If Ω_N is the number of such closed paths, we have [9]:

$$\sum_{A=-A_{max}}^{A_{max}} P_N(\frac{A}{N})\exp(\frac{ixA}{N}) = \Omega_N^{-1}\sum_{\Gamma}\exp(\frac{ixA(\Gamma)}{N}) = \Omega_N^{-1}T(H^N)\mid_{\gamma=\frac{x}{N}}, \tag{19}$$

where A_{max} is the maximum algebraic area closed by Γ with nonvanishing probability. Our aim is to find the formula for $P_N(a)$ where a is renormalized area $a = \frac{A}{N}$, and to investigate its behaviour.
From (7), (8) and (16) we obtain:

$$H = w(\vec{a}) + w(\vec{b}) + w(\vec{-a}) + w(\vec{-b}) = 2i(q-q^{-1})j_y. \tag{20}$$

Thus we observe that the Harper Hamiltonian operator is a generator of the $su_q(2)$ algebra. If the eigenvalue of j_3 is m, the eigenvalue of j_z will be $\pm q^m$. Using (17) or its equivalent :

$$T(w(c)) = \delta_{c,0}. \tag{21}$$

one can show that:

$$T(w(\vec{b}-\vec{a})) = T(\sum_a w(\vec{a})) = T(j_y) = 0, \tag{22}$$

This result also follows from (9) and (10) directly. A straightforward calculation show that the eigenvalues of H have exponential form(see for example equ.(18)). To obtain the eigenvalues of H,we note that:
1. E depends on the quantum number m, and is a function of q-deformation parameter γ(actualy γ and N are not independent, $\gamma = \frac{x}{N}$, see equ.(19), and E can be a function of both of them). It is also has exponential form (for more details see appendix A) and satisfies in equ.(22).
From mathematical point of view, if a series of functions

$$\sum_{n=1}^{\infty} f_n(x). \tag{23}$$

is convergent for any x in some interval, then this series defines a function of variable x, $f(x) = \sum_{n=1}^{\infty} f_n(x)$ in this interval. The importance of a series (e.g. a solution of some differential equations) lies in its convergence. For example, if ℓ be an integer then the series expansions of the solution of the Legendre equation $P_\ell(x)$ and $Q_\ell(x)$ are convergent, and they convert to polynomials. The same statement is also true for the hypergeometric functions and the other polynomials.

We know that the energy of a physical system must be finite, therefore the series (6) in [9] is convergent and is a function of γ (or N) which we will try to determine it.

On the other hand we know that the operators j_y and j_z of $su(2)$ algebra do not commute and can not have common eigenvectors, but their eigenvalues are the same. After deformation the eigenvalues of the j_y and j_z will be not the same, but still their eigenvalues will be discrete labelled by integer quantum number m.

2. We know the eigenvalues of H, in the limit of large N [9].

Therefore we can write :

$$E_m = \pm e^{u_m(\gamma)}. \tag{24}$$

If we choose $u_m(\gamma) = -m\frac{\gamma}{4} + h(\gamma)$, then we have:

$$E_m = \pm f(\gamma) e^{-m\frac{\gamma}{4}}, \tag{25}$$

where $f(\gamma) = \exp(h(\gamma))$. As the candidate for $f(\gamma)$ we take :

$$f(\gamma) = 2(e^{i(\frac{\frac{\gamma}{4}}{\sinh(\frac{N\gamma}{4})})} + e^{-i(\frac{\frac{\gamma}{4}}{\sinh(\frac{N\gamma}{4})})}). \tag{26}$$

We show that its $N \to \infty$ is fully consistent with the result of [9]. We tested it numerically(see below). Then we will have:

$$E_m = \pm 4\cos(\frac{\frac{\gamma}{4}}{\sinh(\frac{N\gamma}{4})}) e^{-\frac{m\gamma}{4}}, \tag{27}$$

By introducing the new variable ℓ

$$m = 2\ell + 1, \tag{28}$$

We have:

$$E_\ell = \pm 4\cos(\frac{\frac{\gamma}{4}}{\sinh(\frac{N\gamma}{4})}) e^{-\frac{(2\ell+1)\gamma}{4}}. \tag{29}$$

Expanding E_ℓ for small values of γ, $\gamma \to 0$, we obtain:

$$E_\ell = \pm 4[(1 - \frac{1}{2N^2}) - (2\ell+1)\frac{\gamma}{4} + [(2\ell+1)^2 + 1]\frac{\gamma^2}{32} - O(\gamma^3)], \tag{30}$$

The term $\frac{1}{2N^2}$ can be omitted, because it tends to zero when $N \to \infty$ ($\gamma \to 0$), and equ.(30) reproduces the eigenvalues used by Bellissard[9]. Now $T(H^N)$ is given by:

$$T(H^N) = \sum_{\pm} \sum_{\ell} (E_\ell^{\pm}(\gamma))^N (\frac{\gamma}{2\pi}), \tag{31}$$

where $\gamma = \frac{x}{N}$ is the multiplicity per unit area. This leads to:

$$T(H^N) = \frac{4^{N+1}}{2\pi N} \frac{\frac{x}{4}}{\sinh(\frac{x}{4})} \cos^N(\frac{\frac{x}{4}}{N \sinh(\frac{x}{4})}), \tag{32}$$

for even N. This gives us the exact characteristic function for the distribution of area. In the limit of large N we obtain:

$$T(H^N(x)) = \frac{4^{N+1}}{2\pi N} \frac{\frac{x}{4}}{\sinh(\frac{x}{4})} [1 - \frac{1}{2N} \frac{(\frac{x}{4})^2}{\sinh^2(\frac{x}{4})} + O(\frac{1}{N^2})]. \tag{33}$$

which is also derived by Bellissard [9]. From (19) and (32) we have:

$$\Omega_N = T(H^N) \mid_{\gamma=0} = \frac{4^{N+1}}{2\pi N} \cos^N(\frac{1}{N}), \tag{34}$$

This gives the total number of closed paths of length N. In the limit of large N we have:

$$\Omega_N = \frac{4^{N+1}}{2\pi N}(1 + (O\frac{1}{N})). \tag{35}$$

Using (19), (32), (34) and the normalization condition we can obtain a series expansion for the probability distribution:

$$P_N(a) = \frac{4}{\pi N} \sum_{m=0}^{\infty} \sum_{k=0}^{\infty} \sum_{\ell=0}^{m+1}$$

$$(-1)^\ell G_m^N 2^{2m+1} \binom{2m+k}{k} \binom{2m+2}{2\ell} \frac{(2m+2k+1)^{2m+2-2\ell}(16a^2)^\ell}{((2m+2k+1)^2 + 16a^2)^{2m+2}} \Gamma(2m+2), \tag{36}$$

We have plotted $P_N(a)$ as given by equ.(36) against exact enumeration[11] in fig1. In the summation of equ.(36) we have allowed m to range from 0 to 2 and k from 0 to 7000 for fig.1.(a) and from 0 to 60000 for fig.1.(b) and (c). Comparing fig.1.(a) and (b), we find that a better result will be obtain if we increase the range of summation over k. The same statement is also true for m. $m = 0$ and $m = 1$ give the equ.(9) in [9] and the $\frac{1}{N}$ correction term made by them respectively. Some of the G_m^N coefficients are presented in Appendix B.
In conclusion we observe that the expression presented for the spectrum of H using quantum groups is in good agreement with the results for $N \to \infty$ and also with exact enumeration. If there is other candidate for $f(\gamma)$, which full fills all the requirements, then the difference between it and our expression will be

negligibly small.

Acknowledgment. One of the authors(S.A.A.) would like to thank Professor S. Rouhani for valuable discussions. He is also very grateful to Professor P. Kulish and Professor P. Prešnajder for their careful reading of the manuscript and for their valuable comments and finally I wish to thank K. Khabbazi and O. Naser Ghodsi for their assistance on computational aspects of this work.

Appendix A. The Harper Hamiltonian is:

$$H = e^{a_1} + e^{a_2} + e^{-a_1} + e^{-a_2} = \sum_{a_i} e^{a_i}, \tag{37}$$

where $[a_i, a_j] = u_{ij}(\gamma)$, $e^{a_i} e^{a_j} = e^{a_i + a_j} e^{\frac{1}{2}[a_i, a_j]}$ and $T(e^{a_i}) = \delta_{a_i,0}$. then:

$$H^N = \sum_{a_i} \sum_{a_j} .... \sum_{a_N} e^{a_i} e^{a_j} .... e^{a_N}. \tag{38}$$

therefore we have:

$$T(H^N) = \sum_m e^{u_m(\gamma)}. \tag{39}$$

$u_m(\gamma)$ must be determined using boundary conditions.

Appendix B. The first four G_m^N coefficients are as follows:

$$G_0^N = 1 \tag{40}$$

$$G_1^N = -\frac{1}{2N}. \tag{41}$$

$$G_2^N = \frac{1}{24} \frac{1}{N^3} + \frac{1}{8} \frac{N-1}{N^3}. \tag{42}$$

$$G_3^N = -\frac{1}{720} \frac{1}{N^5} - \frac{1}{48} \frac{N-1}{N^5} - \frac{1}{48} \frac{(N-1)(N-2)}{N^5}. \tag{43}$$

Figure captions.

Fig.1. The exact and approximate distribution for areas of closed random paths. The exact distribution is plotted for $N = 16$. The solid points were obtained through exact enumerations [11].

References.

1. M.Chaichian and A.Demichev, Introduction to quantum groups, pp. 183-185, World Scientific, Singapore (1996).
2. Haru-Tada Sato, Modern Physics Letters A, Vol.9, No.20, 1819-1825, 1994.

3. Haru-Tada Sato, Modern Physics Letters A, Vol.9, No.5, 451-458, 1994.
4. Alimohammadi and A. Shafei Deh Abad, J. Phys. A: Math. Gen. 29, 559-563, 1996.
5. P. Kulish and Yu. Reshtikhin, J. Sov. Math. 23 (1983) 2435 (translation from: Zapiski Nauch. Seminarov LOMI 101 (1981) 101.)
6. G. Drinfeld in Proc. ICM Berkeley, CA, ed A. M. Gleason(Providence, RI:AMS) P 798 1986.
7.M. Jimbo Lett. Math. Phys 10 63, 1985; Lett. Math. Phys 11 247, 1986.
8. M. Chaichian and P. Kulish, Phys. Lett. B 234 (1990) 72.
9. Jean Bellissard et al J. Phys. A: Math. Gen. 30(1997) L707-L709
10. Harper P G 1955 Proc. Phys. Soc. Lond. A 68 874.
Harper P G 1955 Proc. Phys. Soc. Lond. A 68 879.
11. R. Afsari, N. Sadeghi and S. Rouhani, Exact enumeration of closed random path using TMS320 C6201, Preprint No: IS-00-21. physics/0102016.


a
.1
.1e-1
.1e-2
.1e-3
1e-05
-1.4
-1
-0.8
-0.4
0
0.2
0.4
0.6
0.8
1
1.2
1.4


b
.1
.1e-1
.1e-2
.1e-3
1e-05
1e-06
-1.4
-1
-0.8
-0.4
0
0.2
0.4
0.6
0.8
1
1.2
1.4


c
.1
.1e-1
.1e-2
.1e-3
1e-05
1e-06
-1.4
-1
-0.8
-0.4
0
0.2
0.4
0.6
0.8
1
1.2
1.4